

# IX. Electroweak unification

## The problem of divergence

❖ A theory of weak interactions only by means of  $W^\pm$  bosons leads to **infinities**




Figure 108: Examples of divergent processes.

➔ Introduction of the  $Z^0$  boson fixes the problem because the addition of new diagrams **cancel** out the **divergencies**:




Figure 109: Additional processes which cancel the divergence.

# REMINDER:




Figure 110: The basic W lepton and quark vertices (if the third generation is not taken into account).




Figure 111: The basic  $Z^0$  lepton and quark vertices.




Basic vertices with W bosons have:

- Conserved lepton numbers
- Not conserved quark flavour (quark mixing)



Basic vertices with  $Z^0$  bosons have:

- Conserved lepton numbers
- Conserved flavour (no quark mixing)



## Test of flavour conservation

**Flavour** is **conserved** at a  $Z^0$  vertex (in contrast to a W vertex). This can be verified by experiments.

Consider the following two possible processes that change strangeness:

$$K^+ \rightarrow \pi^0 + \mu^+ + \nu_\mu \quad (\text{a})$$

and

$$K^+ \rightarrow \pi^+ + \nu_l + \bar{\nu}_l \quad (\text{b})$$




Figure 112: Decay (a) is allowed; decay (b) – forbidden

The measured upper limit on the ratio of the decay rates (b) to (a) was until recently:

$$\frac{\sum_l \Gamma(K^+ \rightarrow \pi^+ + \nu_l + \bar{\nu}_l)}{\Gamma(K^+ \rightarrow \pi^0 + \mu^+ + \nu_\mu)} < 10^{-7}$$

# E787 - A rare kaon decay experiment




Figure 113: Side and front views of one half of the E787 experiment.

- ❖ The BNL experiment **E787** is a **fixed target experiment** that uses a  $K^+$  beam created by 24 GeV protons from the AGS accelerator.
- The **Kaons are stopped** in a target made of scintillating fibers and the decay of the  $K^+$  at rest is then studied.
- The **momentum, energy and range** of the particle from the decay is measured.
- **Two candidate events** for  $K^+ \rightarrow \pi^+ + \nu_l + \bar{\nu}_l$  have been found after many years of running.




Figure 114: One of the two rare Kaon decay events found by the E787 experiment.

→ The result of the measurement was:

$$\frac{\sum_l \Gamma(K^+ \rightarrow \pi^+ + \nu_l + \bar{\nu}_l)}{\Gamma(K^+ \rightarrow \pi^0 + \mu^+ + \nu_\mu)} = \frac{1,6 \times 10^{-10}}{0,033} = 5 \times 10^{-9}$$

→ The two events could be explained by second-order charged current interactions:




Figure 115: Feynman diagrams of higher-order charged current interactions resulting in  $K^+ \rightarrow \pi^+ + \nu_l + \bar{\nu}_l$

→ Due to the t-d vertex in the third diagram above it was also possible to set limits on the  $V_{td}$  element in the **Cabibbo-Kobayashi-Maskawa matrix**:

$$0,007 < |V_{td}| < 0,030$$

## The unification condition and masses

The coupling constants at  $\gamma$ -,  $W^\pm$ - and  $Z^0$ -vertices are not independent from each other. In order for all **infinities to cancel** in electroweak theory, the **unification relation** and the **anomaly condition** have to be fulfilled.

→ The *unification condition* establishes a relation between the electroweak coupling constants:

$$\sqrt{\frac{\pi \cdot \alpha_{em}}{2}} = g_W \sin \theta_W = g_Z \cos \theta_W \quad (114)$$

$\theta_W$  is the *weak mixing angle*, or *Weinberg angle*:

$$\cos \theta_W = \frac{M_W}{M_Z} \quad (115)$$

→ The *anomaly condition* relates electric

charges: 
$$\sum_l Q_l + 3 \sum_q Q_q = 0$$

where the factor 3 comes from the number of colors.



Historically, the **W and Z masses** were predicted from **low energy interactions**.




Figure 116: The low energy zero range approximation.

In the zero-range approximation i.e. in the low-energy limit, the charged current reactions are characterized by the **Fermi constant** ( $G_F$ ):

$$\frac{G_F}{\sqrt{2}} = \frac{g_W^2}{M_W^2}$$

From this expression, the unification condition and the definition of  $\theta_W$  one then obtains:

$$M_W^2 = \frac{g_W^2 \sqrt{2}}{G_F} = \frac{\pi \alpha_{em}}{\sqrt{2} G_F \sin^2 \theta_W}$$

$$M_Z^2 = \frac{\pi \alpha_{em}}{\sqrt{2} G_F \cos^2 \theta_W \sin^2 \theta_W}$$

Introducing the **neutral current coupling** constant ( $G_Z$ ) (also in the low energy zero-range approximation) one gets

$$\frac{G_Z}{\sqrt{2}} = \frac{g_Z^2}{M_Z^2}$$

and the weak mixing angle can be expressed as

$$\frac{G_Z}{G_F} = \frac{\sqrt{2} g_Z^2 M_W^2}{\sqrt{2} g_W^2 M_Z^2} = \frac{\frac{\pi\alpha}{2 \cos^2 \theta_W}}{\frac{\pi\alpha}{2 \sin^2 \theta_W}} \cos^2 \theta_W = \sin^2 \theta_W$$

From the **measurements at low energy** of rates of charged and neutral currents reactions ( $G_Z$  and  $G_F$ ) it is therefore possible to determine that:

$$\sin^2 \theta_W = 0,277 \pm 0,014$$

from this measurement at low energies (below the W and Z masses) it was possible to **predict the masses** of W and Z:

$$M_W = 78,3 \pm 2,4 \text{ GeV}/c^2; M_Z = 89,0 \pm 2,0 \text{ GeV}/c^2$$

When the W and Z boson were discovered at CERN with the masses predicted from low energy experiments it was a strong **confirmation** that the **electroweak theory** was correct.

Today the most precise estimation of the Weinberg angle using many measurements give:

$$\sin^2 \theta_W = 0,2255 \pm 0,0021$$

Putting this value into the previous formulas give

$$M_W = 78.5 \text{ GeV} \text{ and } M_Z = 89.3 \text{ GeV}$$

while the direct measurements of the masses give

$$M_W = 80.4 \text{ GeV} \text{ and } M_Z = 91.2 \text{ GeV}$$




Figure 117: Examples of higher order contributions to inverse muon decay

The difference is due to **higher-order diagrams** which were not included in the previous low-energy formulas.

Since the top-quark is involved in higher order corrections, the measurement of electroweak processes could be used to **predict the top-quark mass** before it was discovered:

$$m_t = 170 \pm 30 \text{ GeV}/c^2$$

The directly measured mass of the top quark at Fermilab by CDF is today

$$m_t = 176 \pm 5 \text{ GeV}/c^2$$

in perfect agreement with the prediction !

## Electroweak reactions

❖ In any process in which a **photon** can be exchanged, a  **$Z^0$**  boson can be **exchanged** as well:




Figure 118:  $Z^0$  and  $\gamma$  couplings to leptons and quarks

**Example:** The reaction  $e^+e^- \rightarrow \mu^+\mu^-$  has two dominant contributions:




Figure 119: Dominant contributions to the  $e^+e^-$  annihilation into muons

With simple dimensional arguments one can estimate the **cross section** for the photon- and Z-exchange process at low energy:

$$\sigma_{\gamma} \approx \frac{\alpha^2}{E_{CM}^2} \quad \sigma_Z \approx G_Z^2 E_{CM}^2$$

Where  $E_{cm}$  is the energy of the colliding electron and positron beams.

- One can conclude that at low energies the **photon exchange** process **dominates**.
- However, at energies  $E_{CM}=M_Z$ , this low-energy approximation fails and the  $Z^0$  peak is described by the **Breit-Wigner** formula:

$$\sigma(E_{CM}) = \frac{M^2}{E_{CM}^2} \left[ \frac{C}{(E_{CM}^2 - M^2)^2 + M^2 \Gamma^2} \right]$$

Where  $M$  is the mass and  $\Gamma$  is the decay width.





Figure 120: The cross sections of  $e^+e^-$  annihilation into  $\mu\mu$

$$\sigma(ee \rightarrow \mu\mu) = \frac{12\pi M_Z^2}{E_{CM}^2} \left[ \frac{\Gamma(Z^0 \rightarrow ee)\Gamma(Z^0 \rightarrow \mu\mu)}{(E_{CM}^2 - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \right]$$

## The number of neutrino families

❖ The Z boson can decay in the following way:



Branching ratio ( B ):      0.10                      0.20                      0.70

Decay width (  $\Gamma$  ):      0.25 GeV              0.50 GeV              1.74 GeV

The lifetime ( $\tau$ ) , the branching ratio (B) and the partial decay width ( $\Gamma$ ) are related by

$$\tau = \frac{B}{\Gamma} \qquad B_{xx} = \frac{\Gamma_{xx}}{\Gamma_Z}$$

$$\tau = \frac{B_Z}{\Gamma_Z} = \frac{B_{had} + B_{ll} + B_{\nu\nu}}{\Gamma_{had} + \Gamma_{ll} + \Gamma_{\nu\nu}}$$

$$\tau = \frac{B_{had}}{\Gamma_{had}} = \frac{B_{ll}}{\Gamma_{ll}} = \frac{B_{\nu\nu}}{\Gamma_{\nu\nu}} = 3 \times 10^{-25} s$$

Note:  $1 GeV^{-1} = 6,582 \times 10^{-25} s$






Figure 121: The leptonic decay of the  $Z^0$  into muons.

The peak can be fitted with the Breit-Wigner formula:

$$\sigma(e^+ e^- \rightarrow X) = \frac{12\pi M_Z^2}{E_{CM}^2} \left[ \frac{\Gamma(Z^0 \rightarrow e^+ e^-) \Gamma(Z^0 \rightarrow X)}{(E_{CM}^2 - M_Z^2)^2 + M_Z^2 \Gamma_Z^2} \right]$$

Here  $\Gamma_Z$  is the total  $Z^0$  decay rate, and  $\Gamma_Z(Z^0 \rightarrow X)$  is the decay rates to the final state  $X$ . The height of the peak (at  $E_{CM}=M_Z$ ) is proportional to the product of the branching ratios:

$$B(Z^0 \rightarrow e^+ e^-) B(Z^0 \rightarrow X) \equiv \frac{\Gamma(Z^0 \rightarrow e^+ e^-) \Gamma(Z^0 \rightarrow X)}{\Gamma_Z^2}$$

The fitted parameters of the  $Z^0$  peak in the leptonic and hadronic decay modes give:

$$M_Z = 91,187 \pm 0,007 \text{ GeV}/c^2$$

$$\Gamma_Z = 2,490 \pm 0,007 \text{ GeV}$$

$$\Gamma(Z^0 \rightarrow \text{hadrons}) = 1,741 \pm 0,006 \text{ GeV}$$

$$\Gamma(Z^0 \rightarrow l^+ l^-) = 0,0838 \pm 0,0003 \text{ GeV}$$

→ The decays  $Z^0 \rightarrow l^+ l^-$  and  $Z^0 \rightarrow \text{hadrons}$  account for only about **80%** of all  $Z^0$  decays

→ The remaining decays are those containing only **neutrinos** in the final state since

$$\Gamma_Z = \Gamma(Z^0 \rightarrow \text{hadrons}) + 3\Gamma(Z^0 \rightarrow l^+ l^-) + N_\nu \Gamma(Z^0 \rightarrow \nu_l \bar{\nu}_l) \quad (117)$$

From the measurement of all other partial widths one can therefore estimate the **partial decay to neutrinos** which cannot be measured directly:

$$N_\nu \Gamma(Z^0 \rightarrow \nu_l \bar{\nu}_l) = 0,498 \pm 0,009 \text{ GeV}$$

The decay rate to neutrino pairs can also be **calculated** from the diagrams shown previously and

this gives  $\Gamma(Z^0 \rightarrow \nu_l \bar{\nu}_l) = 0,166 \text{ GeV}$  which

together with  $N_\nu \Gamma(Z^0 \rightarrow \nu_l \bar{\nu}_l) = 0,498 \text{ GeV}$  gives

$$N_\nu = 2,994 \pm 0,011$$




Figure 122: The decay of the  $Z^0$  to hadrons and theoretical predictions based on different assumptions for the number of neutrino families (N)

- There are **no** explicit **restrictions** on the number of generations in the Standard Model.
- However, the analysis of the  $Z^0$  line shape at LEP shows that there are 3 and only 3 kinds of light neutrinos.
- ❖ If neutrinos are assumed having negligible masses as compared with the  $Z^0$  mass, there must be only **THREE generations** of leptons and quarks within the Standard Model.